



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : D06L 3/02, D06M 16/00, C11D 3/386		A1	(11) International Publication Number: WO 99/57360
			(43) International Publication Date: 11 November 1999 (11.11.99)
(21) International Application Number: PCT/DK99/00236 (22) International Filing Date: 29 April 1999 (29.04.99) (30) Priority Data: 0604/98 1 May 1998 (01.05.98) DK 60/084,352 5 May 1998 (05.05.98) US (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): SCHNEIDER, Palle [DK/DK]; Rydtoften 43, DK-2750 Ballerup (DK). DEUSSEN, Heinz-Josef [DE/DK]; Mosevangen 16, DK-2860 Søborg (DK). (74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: ENHANCERS SUCH AS N-HYDROXYACETANILIDE			
(57) Abstract			
The present invention provides a method for bleaching a dye in solution comprising contacting, in an aqueous solution, the dye with a laccase and an enhancing agent of the formula -NOH-CO-, in particular N-Hydroxyacetanilide.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ENHANCERS SUCH AS N-HYDROXYACETANILIDE**FIELD OF INVENTION**

5 The invention relates to a very effective method of bleaching a dye with a laccase enzyme and an enhancing agent. The invention also relates to a detergent composition.

BACKGROUND ART

10 It has earlier been found that coloured substances leached from dyed fabrics could be bleached by means of a phenol oxidizing enzyme. The use of peroxidases or oxidases for inhibiting dye transfer in this way is described in WO 91/05839. It has also been shown that certain oxidizable substances,
15 described as accelerators, enhancing agents or mediators, e.g., metal ions; phenolic compounds such as 7-hydroxycoumarin, vanillin, and p-hydroxybenzenesulfonate, are able to enhance the enzymatic bleaching reactions (see WO 91/05839).

Other types of enhancing agents have also been disclosed,
20 e.g., phenothiazines, phenoxazines, methylsyringate and acetosyringone (see WO 94/12621, WO 95/01426 and WO 95/01426).


Yet another type of enhancing agents are the aliphatic, cycloaliphatic, heterocyclic and aromatic compounds containing NO, NOH or H-N(R)-OH disclosed in WO 94/29425 and WO 97/48786. In WO
25 94/29425 and WO 97/48786 benzotriazoles are preferred as enhancing agents for bleaching dyes.

It is the object of this invention to provide a selected group of -NOH compounds which, together with laccases, are very effective for bleaching dyes in solution.

SUMMARY OF THE INVENTION


It has now surprisingly been found that a selected group of -NOH compounds performs excellently as enhancing agents when used together with laccases for bleaching dyes in solution.

5 Accordingly, the invention provides a method for bleaching a dye in solution comprising contacting, in an aqueous solution, the dye with a laccase and an enhancing agent of the following formula:



10

in which A is:




15

and B is H, or C1-C4 unbranched alkyl wherein said alkyl may contain ether groups, and R2, R3, R4, R5 and R6 are H, NH₂, COOH, CN, SO₃H, CH₃, COCH₃, NO₂, OCH₃, NR₇R₈, COOR₉, or NOH-CO-R₁₀, wherein R₇, R₈, R₉ and R₁₀ are C1-C2 unbranched alkyl.


20

DETAILED DESCRIPTION OF THE INVENTIONEnhancing agents

The present invention relates to a method for bleaching a dye in solution comprising contacting, in an aqueous solution,
25 the dye with a laccase and an enhancing agent of the following formula:



in which A is:



and B is H, or C1-C4 unbranched alkyl wherein said alkyl may contain ether groups, and R2, R3, R4, R5 and R6 are H, NH₂, COOH, SO₃H, CN, CH₃, COCH₃, NO₂, OCH₃, NR₇R₈, COOR₉, or NOH-CO-R₁₀, wherein R₇, R₈, R₉, and R₁₀ are C1-C2 unbranched alkyl; in particular R₂, R₃, R₄, R₅ and R₆ are H, COOH, SO₃H, CH₃, COCH₃, OCH₃, NR₇R₈, or NOH-CO-R₉, wherein R₇, R₈, and R₉ are C1-C2 unbranched alkyl; especially at least three of R₂, R₃, R₄, R₅ and R₆ should be H.

In a preferred embodiment B is H, or C1-C2 unbranched alkyl, and R₂, R₃, R₄, R₅ and R₆ are H, NH₂, COOH, SO₃H, CN, CH₃, COCH₃, NO₂, OCH₃, NR₇R₈, COOR₉, or NOH-CO-R₁₀, wherein R₇, R₈, R₉, and R₁₀ are C1-C2 unbranched alkyl; in particular R₂, R₃, R₄, R₅ and R₆ are H, COOH, SO₃H, CH₃, COCH₃, OCH₃, NR₇R₈, or NOH-CO-R₉, wherein R₇, R₈, and R₉ are C1-C2 unbranched alkyl; especially at least three of R₂, R₃, R₄, R₅ and R₆ should be H.

In another preferred embodiment B is H, or CH₃, and R₂, R₃, R₄, R₅ and R₆ are H, NH₂, COOH, SO₃H, CN, CH₃, COCH₃, NO₂, OCH₃, NR₇R₈, COOR₉, or NOH-CO-R₁₀, wherein R₇, R₈, R₉, and R₁₀ are C1-C2 unbranched alkyl; in particular R₂, R₃, R₄, R₅ and R₆

are H, COOH, SO₃H, CH₃, COCH₃, OCH₃, NR₇R₈, or NOH-CO-R₉, wherein R₇, R₈, and R₉ are C₁-C₂ unbranched alkyl; especially at least three of R₂, R₃, R₄, R₅ and R₆ should be H.

In a particular preferred embodiment, the enhancing agent is N-hydroxyacetanilide.

The enhancing agent of the invention may be present in concentrations of from 1 to 1000 μ M, preferably of from 5 to 500 μ M, and more preferably of from 10 to 200 μ M.

10 Preparation of Enhancing Agents

The enhancing agents described in the present application may be prepared using methods well known to those skilled in the art; a general procedure is outlined in Organic Syntheses 67, 1989, p. 187-192. Some of the enhancing agents are also commercially available.

N-hydroxyacetanilide was produced as described in Organic Syntheses 67, 1989, p. 187-192.

Enzyme

20 The laccase enzyme of the invention may typically be present in concentrations of from 1 to 10000 μ g enzyme protein per liter aqueous solution, in particular of from 5 to 2000 μ g enzyme protein per liter aqueous solution, especially of from 5 to 1000 μ g enzyme protein per liter aqueous solution.

25 Required molecular oxygen will usually be present in sufficient quantity from the atmosphere. If more O₂ is needed, additional oxygen may be added.

Laccase and Laccase Related Enzymes

30 According to the present invention laccase or laccase related enzymes are the preferred enzymes used together with the

selected group of -NOH compounds described above - when a dye in solution is to be bleached.

In the context of this invention, laccases and laccase related enzymes comprise any laccase enzyme comprised by the enzyme classification (EC 1.10.3.2), any catechol oxidase enzyme comprised by the enzyme classification (EC 1.10.3.1), any bilirubin oxidase enzyme comprised by the enzyme classification (EC 1.3.3.5) or any monophenol monooxygenase enzyme comprised by the enzyme classification (EC 1.14.18.1).

10 The above mentioned enzymes may be derived from plants, bacteria or fungi (including filamentous fungi and yeasts) and suitable examples include a laccase derivable from a strain of *Aspergillus*, *Neurospora*, e.g. *N. crassa*, *Podospora*, *Botrytis*, *Collybia*, *Fomes*, *Lentinus*, *Pleurotus*, *Trametes*, e.g. *T. villosa*
15 and *T. versicolor*, *Rhizoctonia*, e.g. *R. solani*, *Coprinus*, e.g. *C. cinereus*, *C. comatus*, *C. friesii*, and *C. plicatilis*, *Psathyrella*, e.g. *P. condelleana*, *Panaeolus*, e.g. *P. papilionaceus*, *Myceliophthora*, e.g. *M. thermophila*, *Scytalidium*, e.g. *S. thermophilum*, *Polyporus*, e.g. *P. pinsitus*,
20 *Pycnoporus*, e.g. *P. cinnabarinus*, *Phlebia*, e.g. *P. radita* (WO 92/01046), or *Coriolus*, e.g. *C. hirsutus* (JP 2-238885).

A laccase derived from *Coprinus*, *Myceliophthora*, *Polyporus*, *Pycnoporus*, *Scytalidium* or *Rhizoctonia* is preferred; in particular a laccase derived from *Coprinus cinereus*,
25 *Myceliophthora thermophila*, *Polyporus pinsitus*, *Pycnoporus cinnabarinus*, *Scytalidium thermophilum* or *Rhizoctonia solani*.

The laccase or the laccase related enzyme may furthermore be one which is producible by a method comprising cultivating a host cell transformed with a recombinant DNA vector which
30 carries a DNA sequence encoding said laccase as well as DNA sequences encoding functions permitting the expression of the DNA sequence encoding the laccase, in a culture medium under

conditions permitting the expression of the laccase enzyme, and recovering the laccase from the culture.

Determination of Laccase Activity (LACU)

- 5 Laccase activity is determined from the oxidation of syringaldazin under aerobic conditions. The violet colour produced is photometered at 530 nm. The analytical conditions are 19 mM syringaldazin, 23 mM acetate buffer, pH 5.5, 30°C, 1 min. reaction time.
- 10 1 laccase unit (LACU) is the amount of enzyme that catalyses the conversion of 1.0 μ mole syringaldazin per minute at these conditions.

Determination of Laccase Activity (LAMU)

- 15 Laccase activity is determined from the oxidation of syringaldazin under aerobic conditions. The violet colour produced is photometered at 530 nm. The analytical conditions are 19 mM syringaldazin, 23 mM Tris/maleate buffer, pH 7.5, 30°C, 1 min. reaction time.
- 20 1 laccase unit (LAMU) is the amount of enzyme that catalyses the conversion of 1.0 μ mole syringaldazin per minute at these conditions.

Dyes or colorants

- 25 In a preferred embodiment, the method of the invention finds application for bleaching of a textile dye or colorant or textile dyes or colorants in solution.

Colorants and dyes are broad classes of natural and synthetic compounds. The following description of dyes/colorants
30 are not intended to be in any way limiting to the scope of the invention as claimed.

Synthetic textile dyes are typically azo compounds (with one or several azo, or diazenediyl, groups), as exemplified by Acid Red 151, Direct Blue 1, Direct Brown 44, and Orange II, or anthraquinone compounds such as Acid Blue 45.

5 Other structural motifs may occur together with these such as Reactive Blue 19.

Some dyes furthermore carry groups capable of coupling to fabric surfaces (reactive dyes), and some dyes are complexed to metal ions. These modifications will often not influence the
10 applicability of the present invention.

A different structure is the indigo moiety, e.g., the soluble dye indigo carmine.

Other dyes and colorants may be of natural origin or may be synthesized as identical to or resembling natural structures.
15 Examples of categories of coloured substances extractable from vegetable sources are polyphenolic, anthocyanine and carotenoid compounds.

Industrial Applications

20 A specific embodiment of the present invention is provided by household and institutional laundering processes.

In such washing and rinsing processes, dyes and colorants present on fabrics may leach into the washing or rinsing liquor and discoloration of the laundry may result. Bleaching of the
25 coloured compounds in solution by the method of the invention may counteract this undesirable effect.

The present invention may also be used for bleaching stains present on fabric: These stains typically originate from red wine, fruit such as black currant, cherry, strawberry and
30 tomato (in particular ketchup and spaghetti sauce), vegetables such as carrots and beetroot, tea, coffee, spices such as curry and paprika, grass, or ball pens/ink.

In another specific embodiment, dyes leached into process water during textile processing may be bleached by the method of the invention to prevent undesirable deposition.

In another specific embodiment, the method of the invention finds application in treatment of waste water, e.g., waste water from the chemical or pharmaceutical industry, from dye manufacturing, from dye-works, or from the textile industry, the method comprising treatment of the waste water with a laccase in the presence of an enhancing agent of the invention.

10 In the above mentioned processes and in other applications of the invention, the enhancing agent may be added at the beginning of the process or later, in one or several additions.

Detergent Compositions

15 The enhancing agent of the invention may be added to and thus become a component of a detergent composition.

The detergent composition of the invention may for example be formulated as a hand or machine laundry detergent composition including a laundry additive composition suitable for pre-
20 treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations, or be formulated for hand or machine dishwashing operations.

In a specific aspect, the invention provides a detergent
25 additive comprising the enhancing agent of the invention and a laccase. The detergent additive as well as the detergent composition may comprise one or more other enzymes such as a protease, a lipase, a cutinase, an amylase, a carbohydrase, a cellulase, a pectinase, a mannanase, an arabinase, a galactanase, a xylanase,
30 an oxidase, e.g., a laccase, and/or a peroxidase.

In general the properties of the chosen enzyme(s) should be compatible with the selected detergent, (i.e. pH-optimum, com-

patibility with other enzymatic and non-enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

Proteases: Suitable proteases include those of animal, vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. The protease may be a serine protease or a metallo protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived from *Bacillus*, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the *Fusarium* protease described in WO 89/06270 and WO 94/25583.

Examples of useful proteases are the variants described in WO 92/19729, WO 98/20115, WO 98/20116, and WO 98/34946, especially the variants with substitutions in one or more of the following positions: 27, 36, 57, 76, 87, 97, 101, 104, 120, 123, 167, 170, 194, 206, 218, 222, 224, 235 and 274.

Preferred commercially available protease enzymes include Alcalase™, Savinase™, Primase™, Duralase™, Esperase™, and Kannase™ (Novo Nordisk A/S), Maxatase™, Maxacal™, Maxapem™, Properase™, Purafect™, Purafect Oxp™, FN2™, and FN3™ (Genencor International Inc.):

Lipases: Suitable lipases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful lipases include lipases from *Humicola* (synonym *Thermomyces*), e.g. from *H. lanuginosa* (*T. lanuginosus*) as described in EP 258 068 and EP 305 216 or from *H. insolens* as described in WO 96/13580, a *Pseudomonas* lipase, e.g. from *P. alcaligenes* or *P. pseudoalcaligenes* (EP 218 272), *P. cepacia* (EP 331 376), *P. stutzeri* (GB 1,372,034), *P.*

fluorescens, *Pseudomonas* sp. strain SD 705 (WO 95/06720 and WO 96/27002), *P. wisconsinensis* (WO 96/12012), a *Bacillus* lipase, e.g. from *B. subtilis* (Dartois et al. (1993), *Biochemica et Biophysica Acta*, 1131, 253-360), *B. stearrowthermophilus* (JP 5 64/744992) or *B. pumilus* (WO 91/16422).

Other examples are lipase variants such as those described in WO 92/05249, WO 94/01541, EP 407 225, EP 260 105, WO 95/35381, WO 96/00292, WO 95/30744, WO 94/25578, WO 95/14783, WO 95/22615, WO 97/04079 and WO 97/07202.

10 Preferred commercially available lipase enzymes include Lipolase™, Lipolase Ultra™ and LipoPrime™ (Novo Nordisk A/S).

Amylases: Suitable amylases (α and/or β) include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, 15 α -amylases obtained from *Bacillus*, e.g. a special strain of *B. licheniformis*, described in more detail in GB 1,296,839.

Examples of useful amylases are the variants described in WO 94/02597, WO 94/18314, WO 96/23873, and WO 97/43424, especially the variants with substitutions in one or more of the 20 following positions: 15, 23, 105, 106, 124, 128, 133, 154, 156, 181, 188, 190, 197, 202, 208, 209, 243, 264, 304, 305, 391, 408, and 444.

Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™ and BAN™ (Novo Nordisk A/S), Rapidase™ and Purastar™ 25 (from Genencor International Inc.).

Cellulases: Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, 30 *Acremonium*, e.g. the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* dis-

closed in US 4,435,307, US 5,648,263, US 5,691,178, US 5,776,757 and WO 89/09259.

Especially suitable cellulases are the alkaline or neutral cellulases having colour care benefits. Examples of such cellulases are cellulases described in EP 0 495 257, EP 0 531 372, WO 96/11262, WO 96/29397, WO 98/08940. Other examples are cellulase variants such as those described in WO 94/07998, EP 0 531 315, US 5,457,046, US 5,686,593, US 5,763,254, WO 95/24471, WO 98/12307 and PCT/DK98/00299.

Commercially available cellulases include Celluzyme™, and Carezyme™ (Novo Nordisk A/S), Clazinase™, and Puradax HA™ (Genencor International Inc.), and KAC-500(B)™ (Kao Corporation).

Peroxidases/Oxidases: Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinus*, e.g. from *C. cinereus*, and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257.

Commercially available peroxidases include Guardzyme™ (Novo Nordisk A/S).

The detergent enzyme(s) may be included in a detergent composition by adding separate additives containing one or more enzymes, or by adding a combined additive comprising all of these enzymes. A detergent additive of the invention, i.e. a separate additive or a combined additive, can be formulated e.g. as a granulate, a liquid, a slurry, etc. Preferred detergent additive formulations are granulates, in particular non-dusting granulates, liquids, in particular stabilized liquids, or slurries.

Non-dusting granulates may be produced, e.g., as disclosed in US 4,106,991 and 4,661,452 and may optionally be coated by methods known in the art. Examples of waxy coating materials are

poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molar weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

The detergent composition of the invention may be in any convenient form, e.g., a bar, a tablet, a powder, a granule, a paste or a liquid. A liquid detergent may be aqueous, typically containing up to 70 % water and 0-30 % organic solvent, or non-aqueous.

The detergent composition comprises one or more surfactants, which may be non-ionic including semi-polar and/or anionic and/or cationic and/or zwitterionic. The surfactants are typically present at a level of from 0.1% to 60% by weight.

When included therein the detergent will usually contain from about 1% to about 40% of an anionic surfactant such as linear alkylbenzenesulfonate, alpha-olefinsulfonate, alkyl sulfate (fatty alcohol sulfate), alcohol ethoxysulfate, secondary alkane-sulfonate, alpha-sulfo fatty acid methyl ester, alkyl- or alkenylsuccinic acid or soap.

When included therein the detergent will usually contain from about 0.2% to about 40% of a non-ionic surfactant such as alcohol ethoxylate, nonylphenol ethoxylate, alkylpolyglycoside, alkyldimethylamineoxide, ethoxylated fatty acid monoethanolamide,

fatty acid monoethanolamide, polyhydroxy alkyl fatty acid amide, or N-acyl N-alkyl derivatives of glucosamine ("glucamides").

The detergent may contain 0-65 % of a detergent builder or complexing agent such as zeolite, diphosphate, triphosphate, 5 phosphonate, carbonate, citrate, nitrilotriacetic acid, ethylenediaminetetraacetic acid, diethylenetriaminepentaacetic acid, alkyl- or alkenylsuccinic acid, soluble silicates or layered silicates (e.g. SKS-6 from Hoechst).

The detergent may comprise one or more polymers. Examples 10 are carboxymethylcellulose, poly(vinylpyrrolidone), poly(ethylene glycol), poly(vinyl alcohol), poly(vinylpyridine-N-oxide), poly(vinylimidazole), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid copolymers.

15 The detergent may contain a bleaching system which may comprise a H_2O_2 source such as perborate or percarbonate which may be combined with a peracid-forming bleach activator such as tetraacetylenediamine or nonanoyloxybenzenesulfonate. Alternatively, the bleaching system may comprise peroxyacids of e.g. 20 the amide, imide, or sulfone type.

The enzyme(s) of the detergent composition of the invention may be stabilized using conventional stabilizing agents, e.g., a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative, e.g., 25 an aromatic borate ester, or a phenyl boronic acid derivative such as 4-formylphenyl boronic acid, and the composition may be formulated as described in e.g. WO 92/19709 and WO 92/19708.

The detergent may also contain other conventional detergent ingredients such as e.g. fabric conditioners including clays, 30 foam boosters, suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil redeposition agents, dyes, bacteri-

cides, optical brighteners, hydrotropes, tarnish inhibitors, or perfumes.

It is at present contemplated that in the detergent compositions any enzyme may be added in an amount corresponding to 5 0.01-100 mg of enzyme protein per liter of wash liquor, preferably 0.05-5 mg of enzyme protein per liter of wash liquor, in particular 0.1-1 mg of enzyme protein per liter of wash liquor.

The enhancing agent of the invention and a laccase may additionally be incorporated in the detergent formulations disclosed in WO 97/07202 which is hereby incorporated as reference. 10

The following examples further illustrate the present invention, and they are not intended to be in any way limiting to the scope of the invention as claimed.

15

EXAMPLE 1

Bleaching of methyl orange using various laccases and various NOH-type enhancing agents

20 **Materials and methods**

Enzymes

Recombinant *Coprinus cinereus* laccase (rCcL), produced as described in WO 97/08325, dialyzed against borate pH 9, 57.5 LAMU/ml.

25 Recombinant *Myceliophthora thermophila* laccase (rMtL), produced as described in WO 95/33836, 1419 LAMU/ml.

Recombinant *Polyporus pinsitus* laccase, produced as described in WO 96/00290, 1330 LACU/ml.

30 Buffer

Britton-Robinson buffer is composed of:

0.1 M phosphate

0.1 M borate

0.1 M acetate

Dye

5 Methyl orange (Merck, Germany)

Enhancing agents

N-Hydroxyacetanilide

1-Hydroxybenzotriazole

10 2-Hydroxypyridine-N-oxide

N-Benzoyl-N-phenylhydroxylamine

Benzoic acid[6-(methylsulphonyl)indol-1-yl]ester (PFC-202)

Cyanuric acid

3-Hydroxy-1,2,3-benzotriazin-4(3H)-one

15 N-hydroxyphthalimide

2-Ethyl-5-phenylisoxazolium-4'-sulfonate, H₂O

p,p'-Azoxydiphenetole

Quinoline-N-oxide

Iso-quinoline-N-oxide

20 di-2-pyridylketone oxime

Bleaching assay

Final conc.

	50 µl	Methyl orange, AbsU _{464nm} ~ 7	1.4 AbsU
	125 µl	0.1 M Britton-Robinson buffer	50 mM
25	25 µl	2 mM mediator (or water)	0.2 mM
	50 µl	laccase dilution	

Reactions were started by addition of enzyme. Absorbance at 464 nm was followed for 3 minutes (readings every 5 seconds) on a Cobas Fara at 30°C. Initial bleaching rates were determined from the first linear part of the progress curves.

30

Results and discussion

The N-OH compounds listed above were tested as laccase mediators in bleaching of methyl orange with three recombinant laccases (rPpL, rMtL and rCcL) at various pH values. All the kinetic data are summarized in Tables 1-3. Large negative numbers represent the best performance.

It can be seen from the Tables that bleaching with O=C-N-OH types enhancing agents (see N-Hydroxyacetanilide) are surprisingly good compared with the other N-OH types enhancing agents.

	PH 4	pH 5	pH 6	pH 7	pH 8
none (reference)	-0.004	-0.006	-0.011	-0.012	-0.005
N-hydroxyacetanilide	-0.521	-0.719	-0.674	-0.498	-0.136
1-hydroxybenzotriazole	-0.016	-0.018	-0.019	-0.017	-0.005
2-hydroxypyridine-N-oxide	-0.008	-0.013	-0.025	-0.020	-0.007
N-benzoyl-N-phenylhydroxylamine	-0.163	-0.247	-0.529	-0.592	-0.124
PFC-202	-0.005	-0.010	-0.016	0.000	0.011
Cyanuric acid	-0.005	-0.008	-0.012	-0.012	-0.005
3-hydroxy-1,2,3-benzotriazin-4(3H)-one	0.004	-0.004	-0.011	-0.008	-0.002
N-hydroxyphthalimide	-0.006	-0.007	-0.014	-0.011	-0.004
2-ethyl-5-phenylisoxazolium-4'-sulfonate, H ₂ O	0.005	0.000	-0.006	-0.004	-0.001
p,p'-azoxydiphenetole	0.116	0.196	0.002	0.008	0.019
Quinoline-N-oxide	-0.004	-0.006	-0.010	-0.008	-0.004
Iso-quinoline-N-oxide	-0.002	-0.005	-0.008	-0.010	-0.004
di-2-pyridylketone oxime	-0.004	-0.006	-0.008	-0.006	-0.001

Table 1. Initial changes per minute in A464nm of methyl orange using rCcL (rCcL solution diluted 1:200).

	pH 4	pH 5	pH 6	pH 7
none (reference)	-0.018	-0.011	-0.002	-0.002
N-hydroxyacetanilide	-1.306	-1.847	-0.630	-0.101
1-hydroxybenzotriazole	-0.050	-0.019	-0.007	-0.002
2-hydroxypyridine-N-oxide	-0.020	-0.016	-0.010	-0.005
N-benzoyl-N-phenylhydroxylamine	-0.269	-0.296	-0.082	-0.012
PFC-202	-0.020	-0.017	0.001	0.005
Cyanuric acid	-0.013	-0.010	-0.002	0.000
3-hydroxy-1,2,3-benzotriazin-4(3H)-one	-0.016	-0.011	-0.001	0.002
N-hydroxyphthalimide	-0.019	-0.016	-0.005	0.000
2-ethyl-5-phenylisoxazolium-4'-sulfonate, H ₂ O	-0.012	-0.004	-0.002	0.000
p,p'-azoxydiphenetole	0.019	0.086	0.012	0.020
Quinoline-N-oxide	-0.016	-0.011	-0.002	0.000
Iso-quinoline-N-oxide	-0.017	-0.010	-0.004	0.000
di-2-pyridylketone oxime	-0.026	-0.010	-0.001	0.001

Table 2. Initial changes per minute in A464nm of methyl orange using rMtL (rMtL solution diluted 1:300).

	PH 4	pH 5	pH 6	pH 7
none (reference)	-0.073	-0.038	-0.017	-0.002
N-hydroxyacetanilide	-4.743	-6.528	-1.837	-0.055
1-hydroxybenzotriazole	-0.206	-0.077	-0.017	-0.005
2-hydroxypyridine-N-oxide	-0.084	-0.052	-0.024	-0.006
N-benzoyl-N-phenylhydroxylamine	-0.732	-1.402	-0.420	-0.012
PFC-202	-0.047	-0.035	-0.001	0.000
Cyanuric acid	-0.037	-0.022	-0.011	-0.002
3-hydroxy-1,2,3-benzotriazin-4(3H)-one	-0.020	-0.020	-0.008	0.000
N-hydroxyphthalimide	-0.050	-0.035	-0.011	-0.001
2-ethyl-5-phenylisoxazolium-4-sulfonate, H ₂ O	-0.017	-0.017	-0.006	0.001
p,p'-azoxydiphenetole	0.121	0.013	0.017	0.018
Quinoline-N-oxide	-0.074	-0.041	-0.016	-0.002
iso-quinoline-N-oxide	-0.074	-0.037	-0.014	-0.002
di-2-pyridylketone oxime	-0.124	-0.107	-0.053	-0.005

Table 3. Initial changes per minute in A464nm of methyl orange using rPpL (rPpL solution diluted 1:200).

5

EXAMPLE 2

Bleaching of methyl orange with laccase and new -NOH-CO- type mediators

10 Materials and methods

Enzymes

Polyporus pinsitus laccase as described in Example 1.

Dye

Methyl orange (Merck, Germany)

Enhancing agents

5 N-Hydroxyacetanilide

N-(4-cyanophenyl)-N-hydroxyacetamide (CAS 80584-65-2)

Bleaching assay

50 μ l 0.1 mg/ml methyl orange

10 125 μ l 0.1 M Britton-Robinson buffer (see example 1)

25 μ l 2 mM mediator or water (as reference)

50 μ l laccase dilution (1:400)

Methyl orange, buffer and mediator are mixed in a micro-titre plate and the reaction is started by addition of laccase.

15 Absorbance is measured at 450 nm after 15 min.

Results and discussion

N-Hydroxyacetanilide and N-(4-cyanophenyl)-N-hydroxyacetamide were tested as mediators in bleaching of methyl orange with a laccase at pH 4-6. The kinetic data are summarized in Table 4. Low values represent the best performance.

The data show N-(4-cyanophenyl)-N-hydroxyacetamide to be as effective as N-Hydroxyacetanilide at bleaching a dye in solution.

	pH 4	pH 5	pH 6
None (reference)	0.920	1.104	1.183
N-Hydroxyacetanilide	0.077	0.068	0.087
N-(4-cyanophenyl)-N-hydroxyacetamide	0.065	0.055	0.083

25

Table 4. Absorbance measured at 450 nm after 15 minutes.

EXAMPLE 3Bleaching of various dyes with laccase and N-Hydroxyacetanilide**Materials and methods****Enzymes**

Polyporus pinsitus laccase as described in Example 1, except that the concentration is 1726 LACU/ml.

Dyes

- 10 Acid blue 113 (Aldrich)
CSB (Aldrich)
Acid blue 45 (Aldrich)
Cibachron Marine (Ciba Specialty Chemicals)

Enhancing agent

N-Hydroxyacetanilide

Bleaching assay

- 50 μ l dye solution
20 125 μ l 0.1 M Britton-Robinson buffer, pH 5 (see example 1)
25 μ l 2 mM mediator or water (as reference)
50 μ l laccase dilution (1:400) or water (as reference)

Dye, buffer and mediator are mixed in a micro-titre plate and the reaction is started by addition of laccase. Absorbance
25 is measured at 595 nm after 5 min.

Results and discussion

N-Hydroxyacetanilide was used as mediator in bleaching of various dyes with a laccase. The kinetic data are summarized in
30 Table 5. Low values represent the best performance.


The data shows that the mediator enhances bleaching of several different dyes.

Dye	- enzyme - mediator	+ enzyme - mediator	+ enzyme + mediator
Acid blue 113	0.474	0.265	0.062
CSB	0.948	0.459	0.207
Acid blue 45	0.941	0.490	0.123
Cibacron Marine	0.981	0.438	0.235

Table 5. Absorbance measured at 595 nm after 5 minutes.


CLAIMS

1. A method for bleaching a dye in solution comprising contacting, in an aqueous solution, the dye with a laccase and
5 an enhancing agent of the following formula:



in which A is:

10



- and B is H, or C1-C4 unbranched alkyl wherein said alkyl may contain ether groups, and R2, R3, R4, R5 and R6 are H, NH₂,
15 COOH, SO₃H, CN, CH₃, COCH₃, NO₂, OCH₃, NR₇R₈, COOR₉, or NOH-CO-R₁₀, wherein R₇, R₈, R₉ and R₁₀ are C1-C2 unbranched alkyl.

2. A method according to claim 1, in which the enhancing agent is N-hydroxyacetanilide.

20

3. A method according to claim 1, in which the laccase is a microbial laccase.

4. A method according to claim 3, wherein the laccase is derived
25 from *Coprinus*, *Myceliophthora*, *Polyporus*, *Pycnoporus*, *Scytalidium* or *Rhizoctonia*.

5. A method according to claim 4, wherein the laccase is derived from *Coprinus cinereus*, *Myceliophthora thermophila*, *Polyporus pinsitus*, *Pycnoporus cinnabarinus*, *Scytalidium thermophilum* or
5 *Rhizoctonia solani*.


6. A method according to any of claims 1-5, in which said method is a method for inhibiting the transfer of a textile dye from a dyed fabric to another fabric when said fabrics are washed
10 together in a wash liquor.

7. A method according to claim 6, in which the enhancing agent is added at the beginning of, or during the process.


15 8. A method according to claim 1, in which the concentration of the enhancing agent is in the range of from 1 to 1000 μM .

9. A detergent composition comprising a laccase, a surfactant and an enhancing agent of the formula:

20



in which A is:



25

and B is H, or C1-C4 unbranched alkyl wherein said alkyl may contain ether groups, and R2, R3, R4, R5 and R6 are H, NH2, COOH, SO3H, CN, CH3, COCH3, NO2, OCH3, NR7R8, COOR9, or NOH-CO-R10, wherein R7, R8, R9 and R10 are C1-C2 unbranched alkyl.

5

10. A detergent composition according to claim 9, in which the enhancing agent is N-hydroxyacetanilide.

11. A detergent composition according to claim 9, wherein the
10 laccase is derived from *Coprinus*, *Myceliophthora*, *Polyporus*,
Pycnoporus, *Scytalidium* or *Rhizoctonia*.

12. A detergent composition according to claim 11, wherein the
laccase is derived from *Coprinus cinereus*, *Myceliophthora*
15 *thermophila*, *Polyporus pinsitus*, *Pycnoporus cinnabarinus*,
Scytalidium thermophilum or *Rhizoctonia solani*.

13. A detergent composition according to any of claims 9-12,
which further comprises one or more other enzymes, in particular
20 a protease, a lipase, an amylase, a cellulase, and/or a
cutinase.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00236

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: D06L 3/02, D06M 16/00, C11D 3/386

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: D06L, D06M, C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, CAPLUS, EPODOC, JAPIO

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	DE 19723912 A1 (CONSORTIUM FÜR ELEKTROCHEMISCHE INDUSTRIE GMBH), 10 December 1998 (10.12.98) --	1-13
X	EP 0825294 A1 (CONSORTIUM FÜR ELEKTROCHEMISCHE INDUSTRIE GMBH), 25 February 1998 (25.02.98) --	1-13
A	DE 4445088 A1 (IBV INDUSTRIELLE BIOVERFAHREN), 20 June 1996 (20.06.96) --	1-13
A	WO 9105839 A1 (NOVO NORDISK A/S), 2 May 1991 (02.05.91) -- -----	1-13

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

29 June 1999

Date of mailing of the international search report

29 -07- 1999

Name and mailing address of the ISA:

Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. - 46 8 666 02 86

Authorized officer

Anna Sjölund/E1s
Telephone No. + 46 8 782 25 00

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/06/99

International application No.

PCT/DK 99/00236

Patent document cited in search report			Publication date	Patent family member(s)	Publication date
DE	19723912	A1	10/12/98	NONE	
EP	0825294	A1	25/02/98	AU 691450 B	14/05/98
				AU 3325497 A	19/02/98
				BR 9704344 A	19/01/99
				CA 2212318 A	13/02/98
				DE 19632623 A	19/02/98
				JP 2846305 B	13/01/99
				JP 10077589 A	24/03/98
				NO 973704 A	16/02/98
				NZ 314955 A	27/04/98
				PL 321586 A	16/02/98
DE	4445088	A1	20/06/96	NONE	
WO	9105839	A1	02/05/91	AT 87029 T	15/04/93
				AT 108484 T	15/07/94
				AU 646645 B	03/03/94
				AU 6515790 A	16/05/91
				AU 6516090 A	16/05/91
				CA 2067748 A,C	14/04/91
				CN 1051600 A	22/05/91
				DE 69010691 D,T	16/03/95
				DK 495836 T	07/06/93
				DK 497794 T	07/11/94
				EP 0495836 A,B	29/07/92
				SE 0495836 T3	
				EP 0497794 A,B	12/08/92
				SE 0497794 T3	
				ES 2057593 T	16/10/94
				FI 100109 B	00/00/00
				FI 921315 A	26/03/92
				FI 921606 A	10/04/92
				IE 63940 B	28/06/95
				JP 2801398 B	21/09/98
				JP 5500899 T	25/02/93
				JP 5503542 T	10/06/93
				MX 172807 B	13/01/94
				PT 95553 A	14/08/91
				TR 26687 A	15/05/95
				US 5273896 A	28/12/93
				US 5605832 A	25/02/97
				US 5648262 A	15/07/97
				US 5700770 A	23/12/97
				US 5712153 A	27/01/98
				US 5855621 A	05/01/99
				WO 9105858 A	02/05/91